

Lists of Larval Worms from Marine Invertebrates of the Pacific Coast of North America

HILDA LEI CHING

Hydra Enterprises Ltd., P.O. Box 2184, Vancouver, British Columbia, Canada V6B 3V7

ABSTRACT: Immature stages of 73 digenetic trematodes are listed by their families, marine invertebrate hosts, and localities and then cross listed according to their molluscan hosts. The list contains many new host records, and larval stages of 15 digenetic trematodes were newly recorded from gastropods from British Columbia and California. A list of immature stages of 14 cestodes, acanthocephalans, and nematodes is also included.

KEY WORDS: Digenea, Cestoda, Acanthocephala, Nematoda, marine mollusks, invertebrate hosts, Pacific coast, North America.

Although immature stages are recognized to be as important as adults in the descriptions of parasites, few studies focus on larval helminths in marine invertebrates. Stunkard (1983) pointed out the difficulties in conducting studies on marine parasites but the necessity to do so should concern all biologists interested in making inventories of parasites and in conserving biodiversity (Pritchard, 1989). The marine ecosystem has become particularly vulnerable to physical changes which in turn can affect both host and parasite populations. My studies of larval digenetic trematodes began in 1957 at Friday Harbor, Washington. Since then, many parasites were found in marine invertebrates from British Columbia to California. Additional sampling was not possible, or when recent collections were attempted, snails and clams were found to be scarce due to oil pollution, or successive changes in the physical environment, or the use of lead-based paints on boats. In this paper I provide lists of larval stages of digenetic trematodes, cestodes, acanthocephalans, and nematodes along with their hosts and localities.

Methods and Materials

Snails, clams, and other invertebrates have been collected by the author at Friday Harbor, Washington (WA), and Vancouver, British Columbia (BC). Eric Cabot collected subtidal snails at Saturnina Island, BC; W. P. Sousa provided infected *Cerithidea* snails from Bolinas Lagoon, California (CA), and G. M. Ruiz gave specimens from clams from Bodega and San Francisco bays, CA. Mollusks were isolated to find free-swimming parasites and then dissected to obtain larval stages. The parasites were studied alive with the use of vital dyes. For permanent preparations, they were usually fixed in hot seawater, transferred to alcohol-formalin-acetic acid and stored in 70% ethanol. They were stained with Semichon's carmine and mounted in permount. Most of the measurements were made on preserved specimens. Measurements are given in micrometers unless otherwise indicated with the mean in parentheses. Drawings were made with the aid of a camera lucida.

Stages that I have reported for the first time are designated with *mihi*. Host and distribution records not previously published are indicated in the list with asterisks. The names of the molluscan hosts and common names when given are cited according to Abbott (1974) and those of other invertebrates are cited according to Austin (1985). Accession numbers are given for specimens in the United States National Museum Helminthological Collections.


List 1. Larval Digeneans from Marine Invertebrates from the North American Pacific Coast

Class TREMATODA
Subclass DIGENEA
Order STRIGEIFORMES

Family CYATHOCOTYLIDAE

1. *Mesostephanus appendiculatus* (Ciurea, 1916) Lutz, 1935
Stage: Furcocercous cercaria
Location: Sporocyst in mantle
Host: *Cerithidea californica* Haldeman, California
Horn Shell

Localities: Upper Newport Bay, Bolinas Lagoon, CA
Records: Martin (1961, 1972), Ching and Sousa, 1986*
USNM Helm. Coll. No. 81066
2. Cyathocotylid species
Stage: Furcocercous cercaria
Location: Sporocyst in digestive gland
Host: *Cerithidea californica*
Localities: Upper Newport Bay, Bolinas Lagoon, CA



Figures 1–7. Marine digeneans from gastropod hosts. 1. Hemiurioid cercaria from *Alvania compacta*. 2. Enenterid cercaria from *A. compacta*. 3. Metacercaria of *Microphallus* sp. from *A. compacta*. 4. Microphallid cercaria from *Littorina scutulata*. 5. Rencolid cercaria from *Cerithidea californica*. 6. Body of lepopocreadiid cercaria from *Nitidella carinata*. 7. Tail of lepopocreadiid cercaria.

Records: Martin (1972), Ching and Sousa, 1986*
USNM Helm. Coll. No. 81067

Family SCHISTOSOMATIDAE

3. *Austrotilharzia variglandis* (Miller and Northup, 1926) Penner, 1953

Stage: Furcocercous cercaria

Location: Sporocyst in gonads

Host: *Ilyanassa obsoleta* (Say), Eastern Mud Nassa

Locality: Vancouver, BC

Record: Ching, 1964*, identification according to Stunkard and Hinchliffe (1952)

4. *Austrotilharzia* sp.

Stage: Furcocercous cercaria

Location: Sporocyst in digestive gland

Host: *Cerithidea californica*

Locality: Upper Newport Bay, CA

Record: Martin (1972)

5. *Cercaria littorinalinae* Penner, 1950
 Stage: Furcocercous cercaria
 Location: Not mentioned, sporocyst in digestive gland?
 Host: *Littorina planaxis* Philippi, Eroded Periwinkle
 Locality: Southern California
 Record: Penner (1950)

Order HEMIURIFORMES

Family HEMIURIDAE

6. *Cercaria B* Miller, 1925
 Stage: Cystophorous cercaria
 Location: Redia in digestive gland
 Host: *Nucella lamellosa* (Gmelin), Frilled Dogwinkle
 Locality: Friday Harbor, WA
 Records: Miller (1925), Ching (1960)
7. *Cercaria C* Ching, 1960
 Stage: Cystophorous cercaria
 Location: Redia in digestive gland
 Host: *Dentalium dalli* Pilsbry and Sharp
 Locality: Friday Harbor, WA
 Record: Ching (1960)
8. Hemiurid Metacercaria
 Stage: Metacercaria
 Location: Hemocoel
 Host: *Mesamphiascus parvus*
 Locality: Friday Harbor, WA
 Record: Ching (1960). The generic name of the host is not listed by Austin (1985), and the copepod may be *Amphiaccus parvus*.
9. Hemiurid Metacercaria
 Stage: Metacercaria
 Location: Unknown
 Host: *Bolinopsis microptera*
 Locality: Puget Sound, WA
 Record: Lloyd (1938). *Bolinopsis microptera* is a synonym of the ctenophore, *B. infundibulum* (Muller), according to Austin (1985).
10. Hemiurid *Cercaria mihi 1* (Fig. 1)
 Stage: Cystophorous cercaria
 Location: Redia in digestive gland
 Host: *Alvania compacta* Carpenter, Compact Alvanina
 Locality: Saturnina I., BC
 Record: Ching, 1977*
 USNM Helm. Coll. No. 81068
 Notes: The prevalence was 3 of 487 of the subtidal snails. The rediae are elongate about 1 mm in length. The cercaria has a delivery tube and 3 appendages.
11. *Lecithaster salmonis* Yamaguti, 1934
 Stage: Cystophorous cercaria
 Location: Redia in digestive gland
 Hosts: *Nucella canaliculata* Duclos, Channeled Dogwinkle; *N. emarginata* (Deshayes), Emarginate Dogwinkle
 Localities and records: Friday Harbor, WA, by Miller as *Cercaria A* (1925) and Ching (1960); BC, by Boyce (1966) as *L. gibbosus*.

Family BUCEPHALIDAE

12. *Cercaria noblei* Giles, 1962
 Stage: Furcocercous cercaria

- Location: Sporocyst in digestive gland
 Host: *Mytilus californianus* Conrad, Californian Mussel
 Locality: Dillon Beach, CA
 Record: Giles (1962)
13. Bucephalid *Cercaria mihi 2*
 Stage: Furcocercous cercaria
 Location: Sporocyst in digestive gland
 Host: *Lyonsia* sp.
 Locality: Friday Harbor, WA
 Record: Ching, 1958*
 Note: The cercaria and host species are similar to those of *Rhipidocotyle* spp. described by Stunkard (1976).

Family FELLODISTOMIDAE

14. *Pronoprymna petrowi* (Layman, 1930) Bray and Gibson, 1980
 Stage: Trichocercous cercaria
 Location: Sporocyst in digestive gland
 Host: *Macoma inconspicua* (Broderip and Sowderby), Inconspicuous Macoma
 Locality: Vancouver, BC
 Records: Ching, 1963*, 1974*, 1989*
 USNM Helm. Coll. No. 81069
 Notes: The prevalence in 1963 was 2.7% or 27 of 1,138 clams. The cercaria resembled young forms found in smelt hosts in the arrangement and development of the gonads (Margolis and Ching, 1965).
15. Fellodistomid Metacercaria *mihi 3*
 Stage: Metacercaria
 Location: Digestive gland
 Host: *Tapes philippinarum* (Adams and Reeve), Japanese Littleneck
 Locality: Vancouver I., BC
 Record: Ching, 1987*
 Notes: About 150 clams from 15 areas were sampled with 1–2 metacercariae found in 7 of 15 clams in 1 area.

Family GYMNOPHALLIDAE

16. *Cercaria margaritensis* Ching, 1982
 Stage: Furcocercous germinal sac
 Location: Germinal sac in extrapallial cavity
 Hosts: *Margarites costalis* (Gould), Northern Rosy Margarite; *M. helacinus* (Phipps), Helicina Margarite; *M. pupillus* (Gould), Puppet Margarite
 Locality: Saturnina I., BC
 Record: Ching (1982)
 USNM Helm. Coll. No. 81070
17. *Gymnophallus deliciosus* (Olsson, 1893) Odhner, 1900
 Stage: Metacercaria
 Location: Extrapallial cavity
 Host: *Mytilus edulis* L., Blue Mussel
 Locality: Vancouver, BC
 Record: Ching, 1963*, identified from Stunkard and Uzzmann (1958)
 USNM Helm. Coll. No. 81071
18. *Gymnophallus somateriae* (Levinson, 1881) Odhner, 1900
 Stage: Furcocercous cercaria
 Location: Sporocyst in digestive gland
 Stage: Metacercaria
 Location: Extrapallial cavity

Host for 2 stages: *Macoma inconspicua*

Locality: Vancouver, BC

Records: Ching (1973), 1963*, 1976* for cercaria
USNM Helm. Coll. Nos. 81072, 81073

19. *Gymnophallus* sp.

Stage: Metacercaria

Location: Extrapallial cavity

Hosts: *Cryptomya californica* (Conrad), California
Glass Mya; *Mya arenaria* (L.), Soft-shell Clam

Record: Ching, 1962*, identified from Stunkard
and Uzmman (1958)

USNM Coll. No. 81074

20. *Lacunovermis conspicuus* Ching, 1965

Stage: Furcocercous cercaria

Location: Sporocyst in digestive gland

Stage: Metacercaria

Host for 2 stages: *Macoma inconspicua*

Locality: Vancouver, BC

Record: Ching (1965a)

21. *Meiogymnophallus multigemmulus* Ching, 1965

Stage: Metacercaria

Location: Sporocyst in digestive gland

Host: *Macoma inconspicua*

Locality: Vancouver, BC

Record: Ching (1965a)

22. *Parvatrema borealis* Stunkard and Uzmman, 1958

Stage: Furcocercous cercaria

Location: Sporocyst in digestive gland

Stage: Metacercaria

Location: Extrapallial cavity

Hosts for 2 stages: *Nuticola tantilla* (Gould), Tan-
tilla Transennella; *Gemma gemma* (Totten),
Amethyst Gem Clam

Localities and records: Friday Harbor, WA, by
Ching, 1959*; San Francisco Bay, CA, by Ogles-
by (1965); Bodega Bay, CA, by Ching and Ruiz,
1986*

USNM Helm. Coll. No. 81075

23. *Parvatrema obscurus* (Ching, 1960) James, 1964

Stage: Metacercaria

Location: Digestive gland

Hosts: "*Collisella*" *scabra* (Gould), Rough Lim-
pet; *Lottia digitalis* (Rathke), Fingered Limpet

Localities: Bodega Bay to San Nicholas I., CA

Record: Ching and Grosholz (1988)

USNM Helm. Coll. No. 79923

24. Gymnophallid Metacercaria

Stage: Metacercaria

Location: Gills, mantle

Host: *Crassostrea gigas* (Thunberg), Giant Pacific
Oyster

Localities: Broughton I., Fanny Bay, Vancouver
I., BC

Record: Bower, 1987, pers. comm., Pacific Bio-
logical Station, Nanaimo, BC

Order PLAGIORCHIIIFORMES

Family ACANTHOCOLPIDAE

25. *Cercaria foliatae* Miller, 1925

Stage: Oculate cercaria

Location: Sporocyst in digestive gland

Host: *Ceratostoma foliatum* (Gmelin), Foliated
Thorn Purpura

Locality: Friday Harbor, WA

Record: Miller (1925). This cercaria is placed in

this family because of its resemblance to the
cercaria of *Stephanostomum baccatum* de-
scribed by Wolfgang (1955) and the close rela-
tionship of the hosts.

Family ENENTERIDAE

26. Enenterid Cercaria *mihi* 4 (Fig. 2)

Stage: Tail-less cercaria

Location: Redia in digestive gland

Host: *Alvania compacta*

Locality: Saturnina I., BC

Record: Ching, 1978*

USNM Helm. Coll. No. 81076

Notes: A single infection was found in 487 snails.

The cercaria appears to be similar to that of
Neophasis lagenformis as described by Koie
(1973). Some opecoelid-like genera with body
spination including *Neophasis* have been placed
in this family by Gibson and Bray (1982).

Family MICROPHALLIDAE

Tribe MARITREMATIDI

27. *Maritrema gratosum* Nicoll, 1907

Stage: Metacercaria

Location: Hemocoel

Host: *Balanus glandula* Darwin

Locality: Vancouver, BC

Record: Ching (1978)

28. *Maritrema loricola* Ching, 1962

Stage: Xiphidiocercaria

Location: Sporocyst in digestive gland

Hosts: *Littorina scutulata* Gould, Checkered Per-
iwinkle; *L. sitkana* Philippi, Sitka Periwinkle

Stage: Metacercaria

Location: Digestive gland

Hosts: *Hemigrapsus nudus* (Dana), *H. oregonensis*
(Dana)

Localities: Vancouver, BC; Bodega Bay, CA

Records: Ching (1962b), Ching, 1986*

29. *Maritrema megametrios* Deblock and Rausch,
1968

Stage: Metacercaria

Location: Hemocoel

Hosts: *Telorchestia traskiana* (Simpson), *Gnori-
mosphaeroma oregonense* Dana

Locality: Vancouver, BC

Record: Ching (1974)

USNM Helm. Coll. No. 81077

30. *Maritrema pacifica* Ching, 1974

Stage: Metacercaria

Location: Hemocoel

Host: *Megalorchestia corniculata* (Stout)

Locality: Santa Barbara, CA

Record: Ching (1974)

USNM Helm. Coll. No. 81078

31. *Probolocoryphe uca* (Sarkisian, 1957) Yamaguti,
1971

Stage: Xiphidiocercaria

Location: Sporocyst in digestive gland

Host: *Cerithidea californica*

Stage: Metacercaria

Location: Muscles

Host: *Uca crenulata* (Lockington)

Localities: Upper Newport Bay, Bolinas Lagoon,
CA

- Records: Maxon and Pequegnat (1949), Sarkisian (1957), Martin (1972) as the larger of 2 stylet cercariae, Sousa (1983) as a microphallid cercaria, Ching and Sousa for cercarial stage, 1986* Cercaria, USNM Helm. Coll. No. 81079
32. *Pseudospilotrema japonicum* Yamaguti, 1939
 Stage: Metacercaria
 Location: Thoracic ganglion
 Host: *Cancer magister* Dana
 Localities: Willapa Bay, WA; Vancouver, BC
 Records: Sparks and Hibbert (1981), identified by Ching, 1988*
- Tribe MICROPHALLIDI
33. *Ascorhytis charadriiformis* (Young, 1949) Ching, 1965
 Stage: Xiphidiocercaria
 Location: Sporocyst in digestive gland
 Hosts: *Littorina scutulata*, *L. sitkana*
 Stage: Metacercaria
 Hosts: *Hemigrapsus oregonensis*, *H. nudus*
 Location: Thoracic ganglion
 Localities: Bodega Bay, CA to Vancouver, BC
 Records: Ching (1962c, 1965b)
34. *Microphallus nicolli* (Cable and Hunninen, 1940) Baer, 1943
 Stage: Xiphidiocercaria
 Location: Sporocyst in digestive gland
 Host: *Olivella plicata* (Sowerby), Purple Dwarf Olive
 Stage: Metacercaria
 Location: Hemocoel
 Host: *Emerita analoga* (Stimpson)
 Locality: Santa Barbara, CA
 Records: Young (1938). Yoshino (1975) reported this species in *Cerithidea californica* but the cercaria in this host is *Probolocoryphe uca* according to Sarkisian (1957).
35. *Microphallus pirum* (Afanassew, 1941) Belopolskaia, 1952
 Stage: Xiphidiocercaria
 Location: Sporocyst in digestive gland
 Host: *Nucella emarginata*
 Stage: Metacercaria
 Location: Abdomen
 Hosts: *Pagurus hirsutiusculus* (Dana), *Telmessus cheiragonus* (Tilsius)
 Locality: Alaska
 Records: Schiller (1954, 1959)
36. *Microphallus pygmaeus* (Levensen, 1881) Baer, 1943
 Stage: Metacercaria
 Location: Sporocyst in digestive gland
 Hosts: *Littorina scutulata*, *L. sitkana*
 Localities: Friday Harbor, WA; Vancouver, BC
 Record: Ching (1961)
 USNM Helm. Coll. No. 81080
37. *Microphallus similis* (Jagerskiold, 1900) Baer, 1943
 Stage: Metacercaria
 Location: Digestive gland
 Host: *Cancer magister*
 Locality: Vancouver, BC
 Record: Ching, 1988*; many specimens from avian hosts have been recovered
38. *Microphallus* sp. *mihi* 5 (Fig. 3)
 Stage: Metacercaria
 Location: Sporocyst in digestive gland
 Host: *Alvania compacta* Carpenter, Compact Alvania
 Locality: Saturnina I., BC
 Record: Ching, 1977*
 USNM Helm. Coll. No. 81081
 Notes: The prevalence was 5.4% or 26 of 487 snails. Measurements on 10 metacercariae: Body length by width, 152–180 by 76–112 (168 by 90). Oral sucker diameter 26–30 (30), ventral sucker diameter 16–20 (18). Pharynx as large as ventral sucker, male papilla diameter, 8–12 (10).
39. Microphallid Cercaria *mihi* 6
 Stage: Xiphidiocercaria
 Location: Sporocyst in digestive gland
 Host: *Lacuna marmorata* Dall
 Locality: Friday Harbor, WA
 Record: Ching, 1959*. Three of 6 snails were infected.
40. Microphallid Cercaria *mihi* 7
 Stage: Xiphidiocercaria
 Location: Sporocyst in digestive gland
 Host: *Trichotropis cancellata* (Hinds), Cancellate Hairy-shell
 Locality: Friday Harbor, WA
 Record: Ching, 1959*. Two of 108 snails were infected.
41. Microphallid Cercaria *mihi* 8 (Fig. 4)
 Stage: Xiphidiocercaria
 Location: Sporocyst in digestive gland
 Host: *Littorina scutulata*
 Locality: Vancouver, BC
 Record: Ching, 1974*
 Notes: I have found only 2 infections in many years of sampling littorine snails. Measurements on 10: Body length by width, 120–171 by 26–42 (132 by 37). Tail short, used for creeping, 26–39 by 5–9 (37 by 7).
- Family OPECOELIDAE
42. *Cercaria searlesiae* Miller, 1925
 Stage: Cotylocercous cercaria
 Location: Redia in digestive gland
 Host: *Searlesia dira* (Reeve)
 Locality: Friday Harbor, WA
 Record: Miller (1925)
43. Metacercaria A Thompson and Margolis, 1987
 Stage: Metacercaria
 Location: Somatic musculature
 Host: *Pandalus jordani* Rathburn, Smooth Pink Shrimp
 Locality: Vancouver I., BC
 Record: Thompson and Margolis (1987)
44. *Neolebouria tinkerbellei* Thompson and Margolis, 1987
 Stage: Metacercaria
 Location: Heart
 Host: *Pandalis jordani*
 Locality: Vancouver I., BC
 Record: Thompson and Margolis (1987)
45. *Podocotyle enophrysi* Park, 1937
 Stage: Cotylocercous cercaria
 Location: Redia in digestive gland

Host: *Lacuna marmorata* Dall

Locality: Saturnina I., BC

Record: Ching (1979)

46. Opecoelid *Cercaria mihi* 9

Stage: Stumpy-tailed cercaria

Location: Sporocyst in digestive gland

Hosts: *Margarites* sp., *M. helacinus*

Localities: Friday Harbor, WA; Saturnina I., BC

Records: Ching, 1957*, 1977*

Note: The cercaria has a short tail with 5–6 glands forming a striated appearance at the tip which is unlike the tails of the cotylocercous cercariae listed above.

47. Opecoelid Metacercaria

Stage: Metacercaria

Location: Hemocoel

Host: *Traskorchestia traskiana*

Locality: Vancouver, BC

Record: Ching, 1977*

48. Opecoelid Metacercaria

Stage: Metacercaria

Location: Abdominal muscles

Host: *Crangon alaskensis* Lockington

Locality: Sinclair Inlet, WA

Record: Morado and Sparks (1983). I have placed this metacercaria in the family Opecoelidae.

Family MONORCHIIDAE

49. *Telolecithus pugetensis* Lloyd and Guberlet, 1932

Stage: Brevifurcate cercaria

Location: Sporocyst in gonads

Host: *Nutricula tantilla* (Gould)

Stage: Metacercaria

Location: Extrapallial cavity

Hosts: *N. tantilla*; *Macoma nasuta*, Conrad Bent-nose Macoma; *Tellina nukuloides* (Reeve), Salmon Tellin

Localities: Yaquina Bay, OR; Friday Harbor, WA

Record: DeMartini and Pratt (1964)

Family RENICOLIDAE

50. *Renicola buehanani* Martin and Gregory, 1951

Stage: Magnacercous cercaria

Location: Sporocyst in mantle

Host: *Cerithidea californica*

Localities: Upper Newport Bay, Bolinas Lagoon, CA

Records: Martin and Gregory (1951), Martin (1955), Yoshino (1975) as *Renicola* sp., Ching and Sousa, 1986*

USNM Helm. Coll. No. 81082

51. *Renicola cerithidicola* Martin, 1971

Stage: Gymnocephalous cercaria

Location: Sporocyst in digestive gland

Host: *Cerithidea californica*

Record: Martin (1971)

52. *Renicola thaidus* Stunkard, 1964

Stage: Xiphidiocercaria

Location: Sporocyst in digestive gland

Host: *Nucella lamellosa*

Stage: Metacercaria

Location: Digestive gland

Host: *Mytilus edulis*

Locality: Vancouver, BC

Records: Ching, 1957*, 1961*, 1969*, Stunkard (1964)

53. Renicolid *Cercaria mihi* 10 (Fig. 5)

Stage: Xiphidiocercaria

Location: Sporocyst in digestive gland

Host: *Cerithidea californica*

Locality: Bolinas Lagoon, CA

Record: Ching and Sousa, 1986*

USNM Helm. Coll. No. 81083

Notes: Sporocyst with yellow pigment, sparsely filled with 2–14 (8) cercariae, 676 by 298 in length and width. Cercarial body finely spined, 179–230 by 58–78 (199 by 69). Suckers equal in size, about 30 in diameter. Short stylet present. Excretory bladder U-shaped with long stem, arms lateral to ventral sucker. Flame cells 2[(3 + 3 + 3) + (3 + 3 + 3)]. Tail length by width, 110–150 by 13–16 (133 by 15).

Order LEPOCREADIIFORMES

Family LEPOCREADIIDAE

54. Lepocreadiid *Cercaria mihi* 11 (Figs. 6, 7)

Stage: Trichocercous cercaria

Location: Redia in digestive gland

Host: *Nitidella carinata* (Hinds), Carinate Dove-shell

Locality: Saturnina I., BC

Record: Ching, 1979*

USNM Helm. Coll. No. 81084

Notes: The prevalence was 5% or 4 of 73 snails.

The cercaria has a trumpet-shaped oral sucker, prominent eye spots, and excretory bladder reaching to the ventral sucker. The tail has 36 pairs of setae.

Order ECHINOSTOMATIFORMES

Family NOTOCOTYLIDAE

55. *Catatropis johnstoni* Martin, 1956

Stage: Oculate, monostome cercaria

Location: Redia in digestive gland

Host: *Cerithidea californica*

Locality: Upper Newport Bay, Bolinas Lagoon, CA

Records: Martin (1955, 1956), Yoshino (1975), Sousa (1983)

56. Notocotylid *Cercaria mihi* 12 (Fig. 8)

Stage: Oculate, monostome cercaria

Location: Redia in digestive gland

Host: *Littorina scutulata*

Locality: Vancouver, BC

Records: Ching (1962a), 1977*

USNM Helm. Coll. No. 81085

Notes: The prevalence was 0.49% or 21 of 4,322 snails sampled in 1977–1978 from 3 locations in Vancouver. Measurements on 10 cercariae: Body length by width, 330–595 by 145–244 (419 by 216). Tail length by width, 376–508 by 33–66 (422 by 43). Three eye spots present.

Family ECHINOSTOMATIDAE


57. *Acanthoparyphium spinulosum* Johnston, 1917

Stage: Cercaria with collar spines

Location: Redia in gonads

Host: *Cerithidea californica*

Stage: Metacercaria



Figures 8–11. Marine digeneans from gastropod hosts. 8. Notocotylid cercaria from *Littorina scutulata*. 9. Echinostomatid cercaria of *Himasthla* sp. from *L. scutulata*. 10. Heterophyid cercaria from *Batillaria zonalis*. 11. Heterophyid cercaria from *Bittium escherichtii*.

- Location: Radular pockets
 Host: *Cerithidea californica*
 Localities: Upper Newport Bay, Bolinas Lagoon, CA
 Records: Martin and Adams (1960, 1961), Yoshino (1975), Sousa (1983), Ching and Sousa, 1986*
 USNM Helm. Coll. No. 81086
58. *Acanthopharyphium* sp.
 Stage: Cercaria with collar spines
 Location: Redia in gonads
 Host: *Cerithidea californica*
 Locality: Upper Newport Bay, CA
 Record: Martin (1972)
59. "*Echinoparyphium*" sp.
 Stage: Cercaria with collar spines
 Location: Redia in gonad, digestive gland

- Host: *Cerithidea californica*
 Localities: Upper Newport Bay, Bolinas Lagoon, CA
 Records: Martin (1972), Sousa (1983), Ching and Sousa, 1986*. Martin (1972) placed this echinostome in this genus, but the 31 spines around the head collar are arranged in a single ring dorsally and laterally, with 2 clustered at each side ventrally.
 USNM Helm. Coll. No. 81087
60. *Himasthla rhigedana* Dietz, 1909
 Stage: Oculate cercaria with collar spines
 Location: Redia in digestive gland
 Host: *Cerithidea californica*
 Localities: Upper Newport Bay, Bolinas Lagoon, CA

Records: Adams and Martin (1960, 1963), Yoshino (1975), Sousa (1983), Ching and Sousa, 1986*
USNM Helm. Coll. No. 81088

61. *Himastha* sp. *Cercaria mihi* 13 (Fig. 9)

Stage: Cercaria with collar spines

Location: Redia in digestive gland

Host: *Littorina scutulata*

Localities: Friday Harbor, WA; Vancouver, BC

Records: Ching (1962a as echinostome I), 1975–1978*, 1988*

USNM Helm. Coll. No. 81089

Notes: The prevalence was 2.5% or 21 of 835 snails at Vancouver and Friday Harbor from 1975 to 1978. At 1 lagoon site at Friday Harbor in 1988, 11% or 23 of 207 snails were infected. Cercarial body spined, with head collar of 29 spines in single continuous row, with 2 ventrally in each corner. Body length by width, 352–454 by 123–217 (400 by 158). Oral sucker 51–59 by 51–56 (57 by 52), ventral sucker, 64–70 by 66–102 (75 by 82). Tail length by width, 237–409 by 33–57 (319 by 41).

62. Echinostomatid Metacercaria

Stage: Metacercaria with collar spines

Location: Gut, mantle, base of gills

Host: *Protothaca staminea* (Conrad), Common Pacific Littleneck

Locality: Sooke, BC

Record: Bower, 1986* pers. comm., Pacific Biological Station, Nanaimo, BC

Family PHILOPHTHALMIDAE

63. *Cloacitrema michiganensis* McIntosh, 1938

Stage: Megalurous cercaria

Host: *Cerithidea californica*

Locality: Upper Newport Bay, CA

Records: Robinson (1952), Martin (1955, 1972), LeFlore et al. (1985)

64. *Parorchis acanthus* (Nicoll, 1906) Nicoll, 1907

Stage: Megalurous cercaria

Location: Redia in digestive gland

Host 1: *Cerithidea californica*

Localities: Upper Newport Bay, Bolinas Lagoon, CA

Records: Martin (1955, 1972), Yoshino (1975), Sousa (1983), Ching and Sousa, 1986*

USNM Helm. Coll. No. 81090

Host 2: *Nucella lamellosa*, *N. emarginata*

Locality: Thetis I., BC

Record: Ching (1978)

USNM Helm. Coll. No. 81091

Family PSILOSTOMIDAE

65. *Psilostomum magniovum* Ching, 1980

Stage: Cercaria with caudal fin fold

Location: Redia in digestive gland

Host: *Littorina scutulata*

Stage: Metacercaria

Location: Extrapallial cavity

Host: *Mytilus edulis* L., Blue Mussel

Locality: Vancouver, BC

Record: Ching (1980)

Order OPISTHORCHIIFORMES

Family HETEROPHYIDAE

66. *Cercaria purpuracauda* Miller, 1925

Stage: Magnacercous cercaria

Location: Redia in digestive gland

Host: *Bittium escherichtii* (Middendorf), Giant Pacific Coast Bittium

Locality: Friday Harbor, WA

Record: Miller (1925)

67. *Cryptocotyle lingua* (Creplin, 1825) Fischeoeder, 1903

Stage: Pleurolophocercous cercaria

Location: Redia in gonads

Host: *Littorina scutulata*

Locality: Vancouver, BC

Record: Ching (1978)

USNM Helm. Coll. No. 81092

68. *Euhaplorchis californiensis* Martin, 1950

Stage: Pleurolophocercous cercaria

Location: Gonad (redial stage in digestive gland)

Host: *Cerithidea californica*

Localities: Upper Newport Bay, Bolinas Lagoon, CA

Records: Martin (1950a, 1955), Yoshino (1975), Sousa (1983), Ching and Sousa, 1986*

USNM Helm. Coll. No. 81093

69. *Parastictodora hancocki* Martin, 1950

Stage: Pleurolophocercous cercaria

Location: Redia in gonads

Host: *Cerithidea californica*

Locality: Upper Newport Bay, CA

Records: Martin (1950b, 1955)

70. *Phocitrema ovale* Martin, 1950

Stage: Pleurolophocercous cercaria

Location: Redia in gonads

Host: *Cerithidea californica*

Localities: Upper Newport Bay, Bolinas Lagoon, CA

Records: Martin (1950c), Ching and Sousa, 1986*
USNM Helm. Coll. No. 81094

71. *Pygidiospoides spindalis* Martin, 1951

Stage: Pleurolophocercous cercaria

Location: Redia in gonads

Host: *Cerithidea californica*

Locality: Upper Newport Bay, CA

Record: Martin (1951)

72. Heterophyid *Cercaria mihi* 14 (Fig. 10)

Stage: Pleurolophocercous cercaria

Location: Redia in gonads

Host: *Batillaria zonalis* (Bruguere) Cumings, False Cerith

Locality: Vancouver, BC

Records: Ching, 1978*, 1989*

USNM Helm. Coll. No. 81095

Notes: The prevalence was 10% or 8 of 77 in 1978 and 22% or 12 of 54 in 1989 in 2 sites in BC. Measurements of 10 cercariae: Body finely-spined, length by width, 139–211 by 66–99 (184 by 80). Tail with finfold ending in small fan around the tip, length by width, 297–475 by 17–36 (366 by 27). Oral sucker with subterminal opening with glandular openings, crenulated walls, 25–46 by 31–37 (31 by 36). Eyespots al-

most square. Penetration glands 14 in total. A pleurolophocercous cercaria reported from this host in California by Whitlatch (1974) may be the same species.

73. Heterophyid *Cercaria mihi* 15 (Fig. 11)

Stage: Pleurolophocercous cercaria

Location: Redia in gonads

Host: *Bittium escherichtii*

Locality: Vancouver, BC

Record: Ching, 1974*

USNM Helm. Coll. No. 81096

Notes: The prevalence was 9.6% or 5 of 52 snails.

Measurements on 9 cercariae: Body length by width, 164–188 by 60–66; tail with short anterior folds, fin folds continuing around bent tip, length by width, 248–409 by 25–30. Oral sucker with 8 spines on the dorsal edge followed by 5 tubular spines in the opening of the mouth. Penetration glands filling body, about 16 in total.

List 2. Molluscan Hosts of Larval Digeneans of the North American Pacific Coast

(Name of mollusc: number of larval digenean in List 1)

GASTROPODA:

Alvania compacta: 10, 26, 38

Batillaria zonalis: 72

Bittium escherichtii: 66, 73

Ceratostoma foliatum: 25

Cerithidea californica: 1, 2, 4, 31, 50, 51, 53, 55, 57, 58, 59, 60, 63, 64, 68, 69, 70, 71

“Collisella” scabra: 23

Ilyanassa obsoleta: 3

Lacuna marmorata: 39, 45

Littorina planaxis: 5

Littorina scutulata: 28, 33, 36, 41, 56, 61, 65, 67

Littorina sitkana: 28, 33, 36

Lottia digitalis: 23

Margarites costalis: 16

Margarites helicinus: 16, 46

Margarites pupillus: 16

Margarites sp.: 46

Nitidella carinata: 54

Nucella canaliculata: 11

Nucella emarginata: 11, 35, 64

Nucella lamellosa: 6, 52, 64

Olivella plicata: 34

Searlesia dira: 42

Trichotropis cancellata: 40

SCAPHOPODA:

Dentalium dalli: 7

BIVALVIA:

Crassostrea gigas: 24

Cryptomya californica: 19

Gemma gemma: 22

Lyonsia sp.: 13

Macoma inconspicua: 14, 18, 20, 21

Macoma nasuta: 49

Mya arenaria: 19

Mytilus californianus: 12

Mytilus edulis: 17, 52, 65

Nuticola tantilla: 22, 49

Protothaca staminea: 62

Tapes philippinarum: 15

Tellina nuculoides: 49

List 3. Larval Tapeworms, Acanthocephalans, and Roundworms from Marine Invertebrates of the North American Pacific Coast

Class CESTODA

Order TETRAPHYLLIDEA

Family PHYLLOBOTHRIIDAE

1. *Anthobothrium* sp.

Stage: Larva

Location: Foot

Hosts: *Tresus nuttalli* (Conrad), Pacific Gaper; *Macoma nasuta*, Bent-nose Macoma; *Protothaca staminea* (Conrad), Common Pacific Littleneck

Locality: Elkhorn Slough, CA

Record: MacGinitie and MacGinitie (1968)

2. *Echeneibothrium* sp.

Stage: Plerocercoid larva

Location: Mantle, foot

Host: *Protothaca staminea*

Locality: Humboldt Bay, CA

Record: Sparks and Chew (1966)

3. *Echeneibothrium* sp.

Stage: Larva

Location: Foot

Host: *Protothaca laciniata*

Locality: Morro Bay, CA

Record: Warner and Katkansky (1969a)

4. *Echeneibothrium* sp.

Stage: Smaller larva

Location: Peripheral and mantle area

Host: *Protothaca staminea*

Locality: Humboldt Bay, CA

Record: Warner and Katkansky (1969b)

5. *Echeneibothrium* sp.

Stage: Larger larva

Location: Viscera

Host: *Protothaca staminea*

Locality: Humboldt Bay, CA

Record: Warner and Katkansky (1969b)

Order TETRARHYNCHIDEA

Family TETRARHYNCHIDAE

6. *Tetrarhynchus* sp. 1 and 2
 Stage: Larvae
 Location: Muscle
 Host: *Dosidicus gigas*
 Locality: Newport Bay, CA
 Record: MacGinitie and MacGinitie (1968)

Order CYCLOPHYLLIDEA

Family HYMENOLEPIDAE

7. *Hymenolepid Cysticercoid mihi* 16
 Stage: Cysticercoid
 Location: Hemocoel
 Host: *Traskorchestia traskiana*
 Locality: Newport Bay, OR
 Record: Ching, 1974*

Phylum ACANTHOCEPHALA

Order PALAEACANTHOCEPHALA

Family POLYMORPHIDAE

1. *Polymorphus kenti* Van Cleave, 1947
 Stage: Cystacanth
 Location: Hemocoel
 Host: *Emerita analoga* (Stimpson)
 Locality: Oregon
 Record: Reish (1950)
2. *Profilicollis botulus* (Van Cleave, 1916) Meyer, 1913
 Stage: Cystacanth
 Location: Rectal ceca
 Host: *Hemigrapsus oregonensis*
 Localities: Bodega Bay, CA to Vancouver, BC
 Record: Ching (1989)
3. Unidentified Cystacanth
 Stage: Cystacanth
 Location: Midgut
 Host: *Paralithodes camtschaticus* (Tilesius)
 Locality: Alaska
 Record: Sparks (1987)

Phylum NEMATODA

Order ASCARIDIDA

Family ANISAKIDAE

1. *Contracaecum* sp.
 Stages: Juvenile and adult
 Location: Hemocoel
 Host: *Pandalis borealis*
 Locality: Open ocean, BC
 Record: Margolis and Butler (1954)

Order SPIRURIDA

Family GNATHOSTOMATIDAE

2. *Echinocephalus pseudouncinatus* Millemann, 1963
 Stage: Juvenile
 Location: Foot
 Host: *Haliotis corrugata* Wood
 Locality: San Clemente I., CA
 Record: Millemann (1963)
 Host: *Centrostephanus coronatus*
 Location: Gonads
 Locality: Southern CA
 Record: Pearse and Timm (1971)
 Location: Adductor muscle

Host: *Argopecten aequisulcatus* (Carpenter)

Locality: Baja California, Mexico

Record: McLean (1983)

Family CYSTIDICOLIDAE

3. *Ascarophis* sp. 1 and 2
 Stage: Juveniles
 Location: Hemocoel
 Hosts: *Pagurus samuelis* (Stimpson), *P. granosimanus* (Stimpson)
 Locality: Bodega Bay, CA
 Record: Poinar and Thomas (1976)
4. *Ascarophis* sp.
 Stage: Juvenile
 Location: Hemocoel
 Hosts: *Pachycheles pubescens* Holmes, *P. rudis*, *Pugettia producta* (Randall), *Hemigrapsus oregonensis*
 Locality: Bodega Bay, CA
 Records: Poinar and Thomas (1976), Poinar and Kuris (1976)

Literature Cited

- Abbott, R. T. 1974. American Seashells, 2nd ed. Van Nostrand Reinhold, New York. 663 pp.
- Adams, J. E., and W. E. Martin. 1960. Life history of *Himasthla* sp., an echinostome trematode. *Journal of Parasitology* 46(Supplement):15.
- , and ———. 1963. Life cycle of *Himasthla rhigedana* Dietz, 1909 (Trematoda: Echinostomatidae). *Transactions of the American Microscopical Society* 82:1-6.
- Austin, W. C. 1985. An Annotated Checklist of Marine Invertebrates in the Cold Temperate Northeast Pacific. 3 vols. Khoyatan Marine Laboratory, British Columbia. 682 pp.
- Boyce, N. 1966. The parasites of central British Columbia pink salmon during their early sea life, with special notes on the trematode *Lecithaster gibbosus*. *Fisheries Research Board of Canada* 26: 813-820.
- Ching, H. L. 1960. Studies on three hemiuroid larvae of Friday Harbor, Washington. *Journal of Parasitology* 46:663-670.
- . 1961. Three trematodes from the harlequin duck. *Canadian Journal of Zoology* 39:373-376.
- . 1962a. Six larval trematodes from the snail, *Littorina scutulata* Gould from San Juan Island, U.S.A., and Vancouver, B.C. *Canadian Journal of Zoology* 40:675-676.
- . 1962b. The description and life cycle of *Maritrema laticola* sp. n. (Trematoda: Microphallidae). *Canadian Journal of Zoology* 41:881-888.
- . 1962c. The life cycle and bionomics of *Levinsoniella charadriiformis* Young, 1949 (Trematoda: Microphallidae). *Canadian Journal of Zoology* 41: 889-899.
- . 1965a. Life cycle of *Lacunovermis conspicuus* n. gen., n. sp. and *Meiogymnophallus multigemmulus* n. gen., n. sp. (Gymnophallidae: Trematoda) from *Macoma inconspicua* and diving ducks from Vancouver, Canada. *Proceedings of the Helminthological Society of Washington* 32:53-63.

- . 1965b. Systematic notes on some North American micropollid trematodes. *Proceedings of the Helminthological Society of Washington* 32: 140–148.
- . 1973. Description of *Gymnophallus somateriae* (Levinson, 1881) from *Macoma inconspicua* and diving ducks from Vancouver, Canada. *Canadian Journal of Zoology* 51:901–906.
- . 1974. Two new species of *Maritrema* (Trematoda: Microphallidae) from the Pacific Coast of North America. *Canadian Journal of Zoology* 52: 865–869.
- . 1978. New marine hosts for *Parorchis acanthus*, *Cryptocotyle lingua*, *Maritrema megametrios*, and *M. gratosum*, trematodes of birds from British Columbia, Canada. *Canadian Journal of Zoology* 56:1877–1879.
- . 1979. The life cycle of *Podocotyle enophrysi* Park, 1937 (Trematoda: Opecoelidae). *Canadian Journal of Zoology* 57:1341–1344.
- . 1980. The life cycle and description of *Psilostomum magnivum* n. sp. (Trematoda: Psilostomidae). *Proceedings of the Helminthological Society of Washington* 47:179–185.
- . 1982. Description of germinal sacs of a gymnophallid trematode, *Cercaria margaritensis* sp. n. in the extrapallial fluid of subtidal snails (*Margarites* spp.) in British Columbia. *Canadian Journal of Zoology* 60:516–520.
- . 1989. *Profilicollis botulus* (Van Cleave, 1916) from diving ducks and shore crabs of British Columbia. *Journal of Parasitology* 75:33–37.
- , and E. Grosholz. 1988. Occurrence of a metacercaria (Trematoda: Gymnophallidae) in acmaeid gastropods, *Lottia digitalis* and “*Collisella*” *scabra*. *Proceedings of the Helminthological Society of Washington* 55:104–105.
- DeMartini, J. D., and I. Pratt. 1964. The life cycle of *Telolecithus pugetensis* Lloyd and Guberlet, 1932 (Trematoda: Monorchidae). *Journal of Parasitology* 50:101–105.
- Gibson, D. I., and R. A. Bray. 1982. A study and reorganization of *Plagioporus* Stafford, 1904 (Digenea: Opecoelidae) and related genera, with special reference to forms from European Atlantic waters. *Journal of Natural History* 16:529–559.
- Giles, D. E. 1962. New bucephalid cercaria from the mussel *Mytilus californianus*. *Journal of Parasitology* 48:293–295.
- Koie, M. 1973. The host-parasite interface and associated structures of the cercaria and adult *Neophasis lagenformis* (Lebour, 1910). *Ophelia* 12: 205–219.
- LeFlore, W. B., H. S. Bass, and W. E. Martin. 1985. The life cycle of *Cloacitrema michiganensis* McIntosh, 1938 (Trematoda: Philophthalmidae). *Journal of Parasitology* 71:28–32.
- Lloyd, L. C. 1938. Some digenetic trematodes from Puget Sound fish. *Journal of Parasitology* 24:117–132.
- MacGinitie, G. E., and N. MacGinitie. 1968. *Natural History of Marine Animals*, 2nd ed. McGraw-Hill, New York. 523 pp.
- Margolis, L., and T. H. Butler. 1954. An unusual and heavy infection of a prawn, *Pandalis borealis* Kroyer, by a nematode, *Contracaecum* sp. *Journal of Parasitology* 40:649–655.
- , and H. L. Ching. 1965. Review of the trematode genera *Bacciger* and *Pentagramma* (Fellodistomatidae) and description of *P. petrowi* (Layman, 1930) n. comb. from marine fishes of the Pacific coast of Canada. *Canadian Journal of Zoology* 43: 381–405.
- Martin, W. E. 1950a. *Euhaplorchis californiensis* n. g., n. sp. Heterophyidae, Trematoda, with notes on its life cycle. *Transactions of the American Microscopical Society* 69:194–209.
- . 1950b. *Parastictodora hancocki* n. gen. n. sp. (Trematoda: Heterophyidae) with observations on its life cycle. *Journal of Parasitology* 36:360–370.
- . 1950c. *Phocitrema ovale* n. g., n. sp. (Trematoda: Opisthorchiidae) with observations on its life cycle. *Journal of Parasitology* 36:552–558.
- . 1951. *Pygidiopsis spindalis* n. gen. n. sp. (Heterophyidae: Trematoda), and its second intermediate host. *Journal of Parasitology* 37:297–300.
- . 1955. Seasonal infections of the snail, *Cerithidea californica* Haldeman with larval trematodes. *Essays in the Natural Sciences in honor of Capt. Allan Hancock*, pp. 203–210. University of California Press, Los Angeles.
- . 1956. The life cycle of *Catantropis johnstoni* n. sp. (Trematoda: Notocotylidae). *Transactions of the American Microscopical Society* 75:117–128.
- . 1961. Life cycle of *Mesostephanus appendiculata* (Ciurea, 1916) Lutz, 1935 (Trematoda: Cyathocotylidae). *Pacific Science* 15:278–281.
- . 1971. Larval stages of renicolid trematodes. *Transactions of the American Microscopical Society* 90:188–194.
- . 1972. An annotated key to the cercariae that develop in the snail *Cerithidea californica*. *Bulletin of the Southern California Academy of Science* 71: 39–43.
- , and J. E. Adams. 1960. Life cycle of *Acanthoparyphium spinulosum* Johnston, 1917 (Echinostomatidae). *Journal of Parasitology* 46(Supplement):35.
- , and ———. 1961. Life cycle of *Acanthoparyphium spinulosum* Johnston, 1917 (Echinostomatidae). *Journal of Parasitology* 47:777–782.
- , and V. Gregory. 1951. *Cercaria buchanani* n. sp., an aggregating marine trematode. *Transactions of the American Microscopical Society* 70: 359–362.
- Maxon, M. G., and W. E. Pequegnat. 1949. Cercariae from upper Newport Bay. *Journal of Entomology and Zoology* 41:30–55.
- McLean, N. 1983. An echinocephalid nematode in the scallop *Argopecten aquiscalcatus* (Mollusca: Bivalvia). *Journal of Invertebrate Pathology* 42: 273–276.
- Millemann, R. E. 1963. Studies on the taxonomy and life history of echinocephalid worms (Nematoda: Spiruroidea) with a complete description of *Echinocephalus pseudouncinatus* Millemann, 1951. *Journal of Parasitology* 49:754–764.
- Miller, H. M. 1925. Larval trematodes of certain

- marine gastropods from Puget Sound. Publications of the Puget Sound Biological Station, University of Washington 5:75-89.
- Morado, J. F., and A. K. Sparks.** 1983. Infection of nervous tissue of shrimp, *Crangon alaskensis*, by trematode metacercariae. *Journal of Invertebrate Pathology* 42:421-423.
- Oglesby, L. C.** 1965. *Parvatrema borealis* (Trematoda) in San Francisco Bay. *Journal of Parasitology* 51:582.
- Pearse, J. S., and R. W. Timm.** 1971. Juvenile nematodes (*Echinocephalus pseudouncinatus*) in the gonads of sea urchins (*Centrostephanus coronatus*) and their effect on host gametogenesis. *Biological Bulletin* 140:95-103.
- Penner, L. R.** 1950. *Cercaria littoralinaliae* sp. nov., a dermatitis-producing schistosome cercaria from *Batillaria minima* (Gmelin). *Journal of Parasitology* 36:466-472.
- Poinar, G. O., Jr., and A. M. Kuris.** 1976. Juvenile *Ascarophis* (Spirurida: Nematoda) parasitizing intertidal decapod Crustacea in California with notes on prevalence and effects on host growth and survival. *Journal of Invertebrate Pathology* 26:375-382.
- , and G. M. Thomas. 1976. Occurrence of *Ascarophis* (Nematoda: Spiruridea) in *Callinassa californiensis* Dana and other decapod crustaceans. *Proceedings of the Helminthological Society of Washington* 43:28-33.
- Pritchard, M. H.** 1989. Biodiversity bill. *American Society of Parasitologists Newsletter* 11:29.
- Reish, D. J.** 1950. Preliminary note in the life cycle of the acanthocephalan, *Polymorphus kenti* Van Cleave, 1947. *Journal of Parasitology* 36:496.
- Robinson, H. W.** 1952. A preliminary report on the life cycle of *Cloacitrema michiganensis* McIntosh, 1938 (Trematoda). *Journal of Parasitology* 38:368.
- Sarkisian, L. N.** 1957. *Maritrema uca*, new species (Trematoda: Microphallidae) from the fiddler crab *Uca crenulata* (Lockington). *Wasman Journal of Biology* 15:35-48.
- Schiller, E. L.** 1954. Studies on the helminth fauna of Alaska XVII. Notes on the intermediate stages of some helminth parasites of the sea otter. *Biological Bulletin* 106:107-121.
- . 1959. Observations on the morphology and life cycle of *Microphallus pirum* (Afanas'ev, 1941). *Transactions of the American Microscopical Society* 78:65-76.
- Sousa, W. P.** 1983. Host life history and the effect of parasitic castration on growth: a field study of *Cerithidea californica* Haldeman (Gastropoda: Prosobranchia) and its trematode parasites. *Journal of Experimental Marine Biology and Ecology* 73:273-296.
- Sparks, A. K.** 1987. The red king crab (*Paralithodes camtschatica*) as a host for juvenile acanthocephalans. *Journal of Invertebrate Pathology* 50:166-168.
- , and K. K. Chew. 1966. Gross infestation of the littleneck clam, *Venerupis staminea* with a larval cestode (*Echeneibothrium* sp.). *Journal of Invertebrate Pathology* 8:413-416.
- , and J. Hibbert. 1981. A trematode metacercaria encysted in the nerves of the Dungeness crab, *Cancer magister*. *Journal of Invertebrate Pathology* 38:88-93.
- Stunkard, H. W.** 1964. Studies on the trematode genus *Renicola*: observations on the life-history, specificity, and systematic position. *Biological Bulletin* 126:467-489.
- . 1976. The life cycles, intermediate hosts, and larval stages of *Rhipidocotyle transversale* Chandler, 1935 and *Rhipidocotyle lintoni* Hopkins, 1954: life cycles and systematics of bucephalid trematodes. *Biological Bulletin* 150:294-317.
- . 1983. The marine cercariae of the Woods Hole, Massachusetts region, a review and a revision. *Biological Bulletin* 164:143-162.
- , and M. C. Hinchliffe. 1952. The morphology and life-history of *Microbilharzia variglandis* (Miller and Northrup, 1926) Stunkard and Hinchliffe, 1951, avian blood flukes whose larvae cause "swimmer's itch" of ocean beaches. *Journal of Parasitology* 38:248-265.
- , and J. R. Uzzmann. 1958. Studies on digenetic trematodes of the genera *Gymnophallus* and *Parvatrema*. *Biological Bulletin* 115:276-302.
- Thompson, A. B., and L. Margolis.** 1987. Description of *Neolebouria tinkerbellei* n. sp. (Trematoda: Digenea: Opecoelidae) from experimental fish hosts, and of metacercaria of *N. tinkerbellei* and an unidentified digenean from *Pandalis jordanii* (Decapoda: Penaeidae) from the Pacific coast of Canada. *Canadian Journal of Zoology* 65:188-193.
- Warner, R. W., and S. C. Katkansky.** 1969a. Infestation of the rough-sided littleneck clam, *Protothaca laciniata* in Morro Bay, California, with larval cestodes (*Echeneibothrium* sp.). *Journal of Invertebrate Pathology* 13:125-128.
- , and ———. 1969b. Infestation of the clam *Protothaca staminea* by two species of tetraphyllidean cestodes (*Echeneibothrium* spp.). *Journal of Invertebrate Pathology* 13:129-133.
- Whitlatch, R. B.** 1974. Studies on the population ecology of the salt marsh gastropod *Batillaria zonalis*. *Veliger* 17:47-55.
- Wolfgang, R. W.** 1955. Studies of the trematode *Stephanostomum baccatum* (Nicoll, 1907). III. Its life cycle. *Canadian Journal of Zoology* 33:113-128.
- Yoshino, T. P.** 1975. A seasonal and histological study of larval Digenea infecting *Cerithidea californica* (Gastropoda: Prosobranchia) from Goleta Slough, Santa Barbara, California. *Veliger* 18:156-161.
- Young, R. T.** 1938. The life history of a trematode (*Levinseniella cruzi*?) from the shore birds, *Limosa fedoa* and *Catoptrophorus semipalmatus inornatus*. *Biological Bulletin* 74:319-329.